



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Reports of States and yearly and monthly reports of cities of the United States.

CONNECTICUT—*New Haven*.—Month of April, 1899. Estimated population, 110,000. Total number of deaths, 152, including diphtheria, 1; enteric fever, 2, and 23 from phthisis pulmonalis.

ILLINOIS—*Chicago*.—During the week ended May 20, 1899, 1 certificate of death was returned to the office of the commissioner of health giving influenza as the chief cause, 14 giving diphtheria as the cause, 69 cases being reported, also 22 giving scarlet fever as the cause, 107 cases being reported.

MICHIGAN.—Reports to the State board of health, Lansing, for the week ended May 20, 1899, from 74 observers, indicate that tonsillitis, pneumonia, and measles decreased in area of prevalence. Phthisis pulmonalis was reported present at 161, measles at 59, scarlet fever at 73, diphtheria at 15, enteric fever at 15, whooping cough at 15, and cerebro-spinal meningitis at 14 places.

MISSOURI—*St. Louis*.—Month of March, 1899. Estimated population, 623,000—white, 587,000; colored, 36,000. Total number of deaths, 986—white, 854; colored, 132, including diphtheria, 21; enteric fever, 4; measles, 1; scarlet fever, 3; smallpox, 2; whooping cough, 10, and 118 from phthisis pulmonalis.

PENNSYLVANIA—*Williamsport*.—Month of April, 1899. Estimated population, 30,000. Total number of deaths, 31, including enteric fever, 2, and 3 from phthisis pulmonalis.

WISCONSIN—*Milwaukee*.—Month of April, 1899. Estimated population, 280,000. Total number of deaths, 326, including diphtheria, 4; enteric fever, 1; measles, 4; scarlet fever, 3; whooping cough, 1, and 39 from phthisis pulmonalis.

MORTALITY TABLE, CITIES OF THE UNITED STATES.

Cities.	Week ended.	Population, U. S. Census of 1890.	Total deaths from all causes.	Deaths from—									
				Phthisis pulmonalis.	Yellow fever.	Smallpox.	Varioloid.	Cholera.	Typhus fever.	Enteric fever.	Scarlet fever.	Diphtheria.	Measles.
Allentown, Pa.....	May 21.....	25,228	6	1
Aurora, Ill.....	May 20.....	19,688	12	2
Baltimore, Md.....	May 27.....	434,439	187	24	1	1
Baton Rouge, La.....	May 13.....	10,478	5
Do.....	May 20.....	10,478	5	1
Binghamton, N. Y.....	May 27.....	35,005	12	1
Brockton, Mass.....	May 13.....	27,294	6	2
Do.....	May 20.....	27,294	6
Camden, N. J.....	do.....	53,313	21	1
Charleston, S. C.....	do.....	a 54,955	b 44	3
Chelsea, Mass.....	do.....	27,909	7
Chicago, Ill.....	May 27.....	1,099,850	440	49	5	7	16	3
Chicopee, Mass.....	May 20.....	14,050	4	2	1
Cleveland, Ohio.....	do.....	261,353	94	2	1	1
Dayton, Ohio.....	May 27.....	61,220	26	4	1

a Estimated population, 65,165—white, 28,870; colored, 36,295.

b White, 12; colored, 32.